

Florida sleeve is a safe and effective technique for valve salvage in acute stanford type A aortic dissection

Mahmoud Alhussaini MD, PhD¹ | Eric I. Jeng MD² | Tomas D. Martin MD² |
Amber Fillion² | Thomas M. Beaver MD, MPH²  | George J. Arnaoutakis MD² 

¹Cardiothoracic Surgery Department, Assiut University, Assiut, Egypt

²Department of Surgery, Division of Cardiovascular Surgery, University of Florida, Gainesville, Florida, USA

Correspondence

George J. Arnaoutakis, MD, Division of Cardiovascular Surgery, University of Florida, PO Box 100129, Gainesville, FL 32610, USA.
Email: george.arnaoutakis@surgery.ufl.edu

Abstract

Objective: Valve-sparing root replacement is commonly used for management of aortic root aneurysms in elective setting, but its technical complexity hinders its broader adoption for acute type-A aortic dissection (ATAAD). The Florida sleeve (FS) procedure is a simplified form of valve sparing aortic root reconstruction that does not require coronary reimplantation. Here, we present our outcomes of the FS repair in patients with dilated roots in the setting of an ATAAD.

Methods: We retrospectively reviewed 24 consecutive patients (2002–2018) treated with FS procedure for ATAAD. Demographic, operative, and postoperative outcomes were queried from our institutional database. Long term follow-up was obtained from clinic visits for local patients, and with telephone and telehealth measures otherwise.

Results: Mean age was 49 ± 14 years with 19 (79%) males. Marfan syndrome was present in 4 (16.7%) patients and 14 (58.3) had $\geq 2+$ aortic insufficiency (AI). Nine (37.2%) had preoperative mal-perfusion or shock. The FS was combined with hemi-arch replacement in 15 (62.5%) patients and a zone-2 arch replacement in 9 (37.5%) patients. There were 2 (8.3%) early postoperative mortalities. Median follow-up period was 46 months (range, 0.3–146). The median survival of the entire cohort was 143.4 months. One patient (4.2%) required redo aortic valve replacement for unrelated aortic valve endocarditis at 30 months postoperatively.

Conclusion: FS is simplified and reproducible valve-sparing root repair. In appropriate patients, it can be applied safely in acute Stanford type-A aortic dissection with excellent early and long-term results.

KEYWORDS

acute aortic dissection, aorta and great vessels, valve repair/replacement

Abbreviations: Ai, aortic insufficiency; AKI, acute kidney injury; ATAAD, acute type A aortic dissection; AVR, aortic valve replacement; COPD, chronic obstructive pulmonary disease; CPB, cardiopulmonary bypass; DHCA, deep hypothermic circulatory arrest; EF, ejection fraction; FS, Florida sleeve; LVEDD, left ventricular end diastolic diameter; LVESD, left ventricular end systolic diameter; TEE, transesophageal echocardiographic; VSRR, valve sparing root replacement.

This study was accepted at the AATS Aortic Symposium, April 23–24, 2020, New York, NY, USA.

1 | BACKGROUND

Valve sparing root replacements (VSRR) since their introduction by Yacoub (remodeling procedure)¹ and David (reimplantation procedure),² have become established alternatives, in experienced hands, to composite graft replacements (Bentall procedure) for aortic root aneurysms.³ Valve sparing procedures avoid the risks of systemic thromboembolism and lifelong anticoagulation associated with mechanical valves and the risks of structural valve deterioration associated with bioprosthetic valves and have excellent clinical outcomes and good long-term durability.^{1,4}

Acute type-A aortic dissection (ATAAD) is a life-threatening emergency and the primary goal of surgery is to save the life of the patient. For dissections involving the aortic root, the Bentall procedure has traditionally been considered the gold standard owing to its technical feasibility and reproducibility.⁵ There are several reports of VSRR techniques in ATAAD with acceptable early and long term results.^{3,6-8} However, the surgical complexity of VSRR and the longer cross-clamp time has slowed wider adoption in the emergent setting.⁹

The Florida sleeve (FS) procedure is a simplified valve sparing root repair that does not require coronary artery reimplantation.¹⁰ The long term durability of this type of repair was published recently yielding excellent results.¹¹ Moreover, the biomechanical characteristics of the FS leads to lower aortic valve stress and prevents possible aortic root distortion or harmful aortic wall stresses.¹² Our aim is to present our outcomes with the FS in ATAAD.

2 | PATIENTS AND METHODS

2.1 | Patients

The University of Florida Institutional Review Board approved this study (#2018-01950). A waiver of informed consent was granted. We retrospectively reviewed 24 consecutive patients at UF Health Aortic Disease Center (The University of Florida College of Medicine, Gainesville, FL) treated with FS procedure for ATAAD from 2002 to 2018. FS was applied by surgeon discretion and indicated when there was an associated root dilatation (aortic root diameter >4.5 and/or >4 cm in patients <60 years or with connective tissue disorder) in the setting of ATAAD.

2.2 | Operative technique

The details of the FS procedure have been described previously.^{10,11,13} Briefly, cardiopulmonary bypass (CPB) was commenced using either axillary, femoral, or central cannulation using the Seldinger technique and under transesophageal echocardiographic (TEE) guidance. After cooling to a core body temperature of 18°C and flat line EEG, deep hypothermic circulatory arrest (DHCA) was initiated with or without retrograde cerebral perfusion. The extent of distal aortic repair was determined. Hemiarch replacement was utilized for intimal tears in the ascending aorta. Zone 2 arch replacements were performed for tears in the arch and proximal descending aorta and/or in younger patients where there was concern for

distal aortic degeneration and high potential for secondary intervention. After de-airing of the distal aortic graft, re-establishing of CPB and during the rewarming phase, the aortic root was addressed. The aortic root was dissected circumferentially to the annular level, and the proximal portions of the coronary arteries were mobilized. To size graft appropriately, a Hegar dilator is used to measure the annulus directly, and a Valsalva graft 6–8 mm larger than the annulus diameter is selected, typically 30 or 32 mm graft for women and 32 or 34 mm graft for men. If additional annular reduction is desired in patients with larger annulus size, an additional fourth annuloplasty suture is placed at nadir of noncoronary cusp. Typically, three subannular 2-0 pledgeted Ethibond (Ethicon US, LLC) mattress sutures are utilized to anchor the FS graft at the root (three in the subcommissural positions, and additional fourth subannular suture at the nadir of the noncoronary cusp) (Figure 1A). Because the coronary arteries are left attached to the aortic root, it can be difficult to place the subannular sutures at the nadir of each cusp, as visualization and needle manipulation in this area is more difficult with the coronary arteries left attached. In contrast, in a David technique where both coronary buttons are detached and then reimplanted, subannular suture placement at the nadir of each cusp is technically more easily accomplished.

With bicuspid valves, subannular sutures were placed under each of the commissures. We were careful with additional subannular sutures in bicuspid valves as they may produce unwanted leaflet elongation, compromising the geometry of coaptation and subsequently valvular competence. While cusp repair can be performed with the FL sleeve technique including in acute dissection settings, we usually perform adjunctive cusp plication techniques in nonemergent scenarios such as aneurysm repair. In this series of acute type A dissections, all of the operations were FL sleeve procedures, but no cusp plications were undertaken. Locations of the coronary arteries were marked on the graft, and vertical slits were made to create coronary keyholes. The subannular anchor sutures were placed through a Valsalva graft (Terumo Aortic Vascutek Ltd.), graft was seated, and the sutures are tied with a Hegar dilator (Jarit Instruments) through the annulus to prevent a purse-string effect. The slits below the coronary arteries were closed using simple sutures making sure these keyholes were large enough to accommodate the coronary arteries. At this time, the Valsalva graft is cut to size and supra-commissural sutures were placed to resuspend the valve, being careful to reapproximate all of the dissected layers of the aorta (Figure 1B). The dissection repair is completed with sandwich of Teflon-felt-aorta-Dacron at the sinotubular junction (STJ) with a running horizontal mattress (Figure 1C). A graft to graft anastomosis of the proximal and distal repairs completes the operation (Figure 1D). Intraoperative TEE was used to assess the quality of valve repair, degree of aortic insufficiency (AI) and ventricular function to detect any segmental wall motion abnormalities that might have happened due to coronary artery impingement.

2.3 | Endpoints and statistical analysis

The primary endpoints chosen were the early mortality and post-operative AI. Additional endpoints included procedural safety,




FIGURE 1 (A) A showing four subannular 2-0 pledgeted mattress sutures at the aortic root (three in the subcommissural positions, and one at the nadir of the left-non coronary cusp). These sutures are anchored to the Valsalva graft. (B) The slits below the coronary arteries were closed using simple sutures. The Valsalva graft is cut at the level of the STJ and three resuspension sutures are paced at each commissure to ensure valve competence. (C) Running horizontal mattress sutures incorporating the Dacron graft, dissected layers of the aorta and Teflon felt strip. (D) Restoring the aortic continuity with graft to graft anastomosis. STJ, sinotubular junction

long-term durability, and freedom from reoperation. Postoperative mortality was defined as mortality at 30 days postoperatively or at time of discharge. Acute kidney injury (AKI) was defined using the AKIN Criteria.¹⁴ We used echocardiography to assess the degree of AI (graded as: 0 = none, 1+ = trace/minimal, 2+ = mild, 3+ = moderate, and 4+ = severe), ejection fraction (EF), left ventricular end diastolic diameter (LVEDD), and left ventricular end systolic diameter (LVESD). Follow-up echocardiography measurements were not available for all patients because some patients had postoperative follow-up echocardiography at outside centers. Long-term follow-up was obtained from clinic visits for local patients, and with telephone and telehealth measures otherwise. Social security death index was utilized for those who could not be contacted.

Continuous variables were presented as the mean \pm SD and categorical data as frequency and percentage. Quantitative measures are expressed as mean \pm 1 SD where appropriate.

Outcome data were analyzed with χ^2 test, Fisher exact test, and the independent Student's *t* test where appropriate. Differences were considered significant if a 95% confidence interval was achieved. Patients' survival rate and freedom from reoperation were evaluated by Kaplan-Meier and life-table methods. All statistics were performed by Graphpad Prism software (Version 8.4; Graphpad software, LLC).

3 | RESULTS

3.1 | Patients demographics and preoperative variables

Twenty-four patients treated with FS procedure for ATAAD, 19 men (79.2%) with mean age of 49 ± 14 years were included in the study.

TABLE 1 Patients demographics and preoperative characteristics

Variables	N = 24 (%)
Age (years) mean \pm SD	49.25 \pm 14.04
Male	19 (79.2)
Body mass index mean \pm SD	28.4 \pm 6.9
Type of dissection (Debakey classification)	
I	21 (87.5)
II	3 (12.5)
Marfan syndrome	4 (16.7)
Bicuspid aortic valve (BAV)	1 (4.2)
Familial aortic aneurysms (non-marfan)	3 (12.5)
Current or ex-smoker	13 (54.5)
Chronic lung disease	3 (12.5)
Diabetes	0 (0)
Hypertension	14 (58.3)
Dyslipidemia	3 (12.5)
Peripheral arterial disease	2 (8.3)
Cerebrovascular disease	2 (8.3)
Chronic renal insufficiency	3 (12.5)
Renal insufficiency on dialysis	1 (4.2)
Arrhythmia	1 (4.2)
Congestive heart failure	1 (4.2)
Previous sternotomy	1 (4.2)
Degree of aortic insufficiency (AI)	
0	5 (20.8)
1+	2 (8.3)
2+	5 (20.8)
3+	3 (12.5)
4+	9 (37.2)
Condition at time of surgery	
Aortic root diameter (mm) mean \pm SD	53.3 \pm 9
Preoperative myocardial infarction	1 (4.2)
Malperfusion syndrome	9 (37.2)
Bloody pericardial effusion	9 (37.5)
Preoperative hematocrit (%) mean \pm SD	35.7 \pm 8.7
Preoperative creatinine (mg/dl) mean \pm SD	1.4 \pm 1.3
Preoperative INR mean \pm SD	1.1 \pm 0.2

Abbreviation: INR, international normalized ratio.

Marfan syndrome was present in 4 (16.7%) patients and 12 (50%) had >2+ AI. Nine (37.2%) had preoperative malperfusion or shock. Mean aortic root diameter was 53.3 \pm 9 mm. The main preoperative features of the patients are summarized in Table 1.

3.2 | Operative variables and postoperative outcomes

Operative and postoperative variables are demonstrated in Table 2. The FS was combined with hemi-arch replacement in 15 (62.5%) patients or arch debranching and zone-2 arch replacement in 9 (37.5%) patients. Mean CPB time was 225.5 \pm 45.8 min and mean myocardial ischemic time was 137.4 \pm 35.5 min. DHCA was used in all patients and this was combined with either antegrade or retrograde cerebral perfusion in 15 (62.5%) patients. Intraoperative TEE showed procedural success in all patients regarding AI and none of the patients had more than mild AI at time of discharge. None of the patients had postoperative neurological complications or required re-exploration for bleeding. There were 2 (8.3%) early postoperative mortalities; one had preoperative renal failure and myocardial infarction and the other had a preoperative ruptured ascending aorta and cardiac tamponade with malperfusion and the postoperative course was complicated by septicemia, renal, and respiratory failure.

3.3 | Follow-up

The median follow-up period was 46 months (range, 0.3–146). Only 1 (4.2%) required redo aortic valve replacement (AVR) for endocarditis (30 months postoperatively) during follow-up (Figure 2). Five (20.8%) patients required intervention on the descending thoracic/thoracoabdominal aorta during follow-up. There were 2 late postoperative mortalities one from respiratory failure complicating COPD 12 years after surgery and the other from sepsis complicating open repair of ruptured thoracoabdominal aneurysm 3 years following ATAAD repair. Echocardiography during follow-up was available in 12 patients at 1–3 years and it showed decrease of mean degree of AI from 2.4 \pm 1.6 preoperatively to 0.9 \pm 1.4 at 1–3 years ($p = .01$) and LVEDD from 54.7 \pm 6.9 mm preoperatively to 46.9 \pm 9.4 mm at 1–3 years ($p = .04$) (Table 3). Freedom from redo 1, 5, and 10 years AVR is estimated to be 100%, 93 \pm 6.9%, and 93 \pm 6.9% respectively. Estimated median survival time is 143.4 months and estimated survival is 85.2 \pm 8% at 1 and 10 years (Figure 3).

4 | DISCUSSION

Although VSRRs are well-established surgical options for management of elective aortic root pathology (chronic dissections or aneurysms), it is not widely performed in the setting of ATAAD with root pathology. VSRR procedures are hindered by their technical complexity and the additional time required to perform the repair and may be not suitable in this emergent clinical setting. Furthermore, there are concerns about the safety and durability and the risk of late valve dysfunction secondary to AI.

The immediate goal in ATAAD is to save the life of the patient and in select cases root replacement with composite valve conduit is considered traditional therapy. Even at centers with extensive VSRR

TABLE 2 Operative variables, postoperative outcomes and follow-up

Variables	N = 24 (5)
Type of arch repair	
Hemi-arch replacement	15 (62.5)
Arch debranching and zone 2 arch replacement	9 (37.5)
Antegrade stent graft deployment (frozen elephant trunk)	1 (4.2)
Concomitant coronary artery bypass grafting (CABG)	2 (8.3)
Florida sleeve Valsalva graft size (mm)	
32	20 (83.3)
34	3 (12.5)
36	1 (4.2)
Cardiopulmonary bypass time (min) mean \pm SD	225.5 \pm 45.8
Myocardial ischemic time (min) mean \pm SD	137.4 \pm 35.5
Arterial cannulation type	
Central cannulation	18 (75)
Femoral cannulation	6 (25)
Circulatory arrest	24 (100)
Cerebral perfusion	15 (62.5)
ACP	7 (29.2)
RCP	7 (29.2)
Combined ACP/RCP	1 (4.2)
Circulatory arrest time (min) mean \pm SD	38.8 \pm 23.5
Cerebral ischemic time (min) mean \pm SD	18.7 \pm 12.1
Postoperative outcomes	
Postoperative degree of aortic insufficiency	
0-1	21 (75)
2	3 (25)
3-4	0 (0)
Postoperative acute kidney injury	5 (20.8)
Postoperative renal failure necessitating dialysis at discharge	3 (12.5)
Arrhythmia	4 (16.7)
Postoperative stroke	0 (0)
Postoperative bleeding necessitating re-exploration	0 (0)
Early postoperative mortality	2 (8.3)
ICU stay (days) mean \pm SD	8.1 \pm 8.4
Postoperative mechanical ventilation time (h) mean \pm SD	114.4 \pm 208
Length of hospital stay (days) mean \pm SD	12 \pm 8.2
>Mild aortic insufficiency in postoperative echocardiography	0 (0)
Follow-up	
Follow-up duration (months) mean \pm SD	53.7 \pm 50.5
Redo aortic valve replacement (AVR)	1 (4.2)
TEVAR	3 (12.5)

(Continues)

TABLE 2 (Continued)

Variables	N = 24 (5)
Open descending/thoracoabdominal aortic repair	2 (8.3)
Estimated survival time (months) mean \pm SEM	125.2 \pm 10.6

Abbreviations: ACP, antegrade cerebral perfusion; ICU, intensive care unit; RCP, retrograde cerebral perfusion; TEVAR, thoracic endovascular aortic repair.




FIGURE 2 Freedom from aortic valve reoperation following Florida sleeve repair for acute type A aortic dissection

experience VSSR in dissection is limited as in Toronto with a quarter of a century of experience there were 28 patients with ATAAD.⁴ The largest experience with David procedure in ATAAD was reported by the Hannover group. They reported 109 David I procedures in ATAAD with 11% operative mortality and 85% freedom from valve-related reoperation at 10 years.¹⁵ The Leipzig group reported 78 VSRRs (51 Yacoub and 27 David) with 30-day mortality of 15%, 8-year survival of 55% and estimated freedom from reoperation 8.9 years.¹⁶ Interestingly, only 2 of their patients (2.6%) had Marfan syndrome (David procedures) and this may explain their preferred use of the Yacoub technique. Recently, Aubin and his colleagues reported their results with 28 David repairs in TAAD with 30-day and 5-year mortality of 18% and 29%, respectively.¹⁷

The FS is a simplified aortic root repair that does not involve replacement of the sinuses of Valsalva. Instead, the entire root unit is enveloped in a Valsalva Dacron graft tailored as a “sleeve,” leaving the coronary artery ostia intact. The horizontal mattress suture line in the setting of acute dissection is used to appose all layers of the aortic wall and accomplishes aortic valve commissural resuspension. A felt strip is used on the inner layer and the dacron graft serves as the external buttress. While this can be performed without, we believe this provides a more hemostatic and robust suture line, especially in the setting of acute dissection. It is important to note that purse-stringing can occur, therefore it is recommended not to overtighten the horizontal mattress suture line. This can be prevented by tying the knot down over an appropriate sized Hegar dilator to avoid over narrowing the STJ.

This technique allows reduction and stabilization of the virtual basal ring and the STJ, while the Valsalva graft allows for normal expansion of the sinuses of Valsalva during cardiac cycle, rendering more physiological biomechanics than the David procedure—with lower leaflet and commissural stress as reported by Gamba and his colleagues.¹² Furthermore, this technique does not require coronary button reimplantation, which can be treacherous if the friable dissection plane extends into either, or both, of the coronary ostia.^{18,19} The main instance for performing David repair in acute dissection in our center would be scenarios where there is significant tear extending down into the sinus segments, precluding external buttress alone and requiring complete root resection for hemostatic purposes. Also there needs to be enough intact aortic tissue above each coronary ostium to perform a hemostatic STJ suture line anastomosis, and if tear encroaches too closely on either coronary ostium, this would be an instance for David repair or modified Bentall operation.

The residual sinus segment is contained within the Valsalva portion of the Dacron graft, and any sinus segment expansion is limited by the external diameters of the Valsalva segment of the graft. We have not identified any de novo cases of dissection in the native sinus segment in patients who have undergone FL Sleeve repair for acute dissection when there was no dissection extending into the root segment at initial presentation. It should be emphasized that FL Sleeve can be performed in instances where dissection extends down into the sinus segment as the layers of the aortic wall are opposed during the horizontal mattress layer at the STJ. Since there are anchoring sutures at the annular level (subannular sutures), and the STJ is maintained by the suture lines, this should be a durable repair and not lead to significant AI. While the current study was undertaken to evaluate short term outcomes to describe the feasibility of this technique in acute dissection, we have not experienced any cases of early recurrent AI, and the long term durability in the setting of acute dissection is an area of current inquiry.


Our group recently published long-term experience with the FS technique with 98% freedom from reoperation and 93% survival at 8 years.¹¹ This is comparable to the long-term experience of David and his colleagues with VSRRs who reported 97% freedom from reoperation and 89% survival at 10 years.⁴

The 8% mortality with FS repair in ATAAD in this series is favorable compared to a recent International Registry of Acute Aortic Dissection (IRAD) report with 18% 30-day mortality in ATAAD.⁹ Procedural success in the form of absence of greater than 2+ degree AI was achieved in all patients. Only one patient required AVR during follow-up but that was due to valve endocarditis suggesting the FS is a stable aortic valve repair for the long term. Echocardiographic data

TABLE 3 Preoperative, early postoperative, and last follow up Echocardiographic data

	Preoperative echocardiography	Early postoperative echocardiography	Last follow-up echocardiography	p value
Degree of AI mean \pm SD	2.4 \pm 1.6	0.3 \pm 0.7	0.9 \pm 1.4	.01
LVEDD (mm) mean \pm SD	54.7 \pm 6.9	47.5 \pm 7.6	46.9 \pm 9.4	.04
LVESD (mm) mean \pm SD	38.5 \pm 5.9	31.9 \pm 7.9	32.1 \pm 6.5	.07
EF (%) mean \pm SD	56.3 \pm 9.6	59.7 \pm 6.9	63.43 \pm 4.2	.08

Abbreviations: AI, aortic insufficiency; EF, ejection fraction; LVEDD, left ventricular end diastolic diameter; LVESD, left ventricular end systolic diameter.

**FIGURE 3** Kaplan-Meier survival curve following Florida sleeve repair for acute type A aortic dissection

showed improvement in LVEDD at 1–3 years which is thought to be the most important indicator of AI.²⁰ Freedom of redo AVR (92.9%) at 10 years and 10 year survival (85.2%) is comparable to other VSRRs in type A dissection.^{15–17}

5 | LIMITATIONS

This study has the following limitations: (1) It is a retrospective single center study of a small group of patients. (2) Follow-up echocardiography was available in only 50% of the study group.

6 | CONCLUSION

The FS is a simplified and reproducible valve-sparing root repair. In carefully selected patients, it can be applied safely in ATAAD with excellent early and long-term results. Further studies are needed comparing larger numbers of patients to composite valved conduit root replacements and other valve sparing root repairs.

ACKNOWLEDGEMENT

This research was funded by Herron Endowed Chair in Cardiothoracic Surgery at The University of Florida.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

ORCID

Thomas M. Beaver  <http://orcid.org/0000-0003-1785-0119>

George J. Arnaoutakis  <http://orcid.org/0000-0002-7648-0747>

REFERENCES

1. Yacoub MH, Gehle P, Chandrasekaran V, Birks EJ, Child A, Radley-Smith R. Late results of a valve-preserving operation in patients with aneurysms of the ascending aorta and root. *J Thorac Cardiovasc Surg.* 1998;115(5):1080-1090.
2. David TE, Feindel CM. An aortic valve-sparing operation for patients with aortic incompetence and aneurysm of the ascending aorta. *J Thorac Cardiovasc Surg.* 1992;103(4):617-621.
3. Ouzounian M, Rao V, Manlihot C, et al. Valve-sparing root replacement compared with composite valve graft procedures in patients with aortic root dilation. *J Am Coll Cardiol.* 2016;68(17):1838-1847.

4. David TE, Feindel CM, David CM, Manlihot C. A quarter of a century of experience with aortic valve-sparing operations. *J Thorac Cardiovasc Surg.* 2014;148(3):872-879.
5. Hagan PG, Nienaber CA, Isselbacher EM, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA.* 2000;283(7):897-903.
6. Lee H, Cho YH, Sung K, et al. Clinical outcomes of valve-sparing root replacement in acute type A aortic dissection. *Scand Cardiovasc J.* 2015; 49(6):331-336.
7. Stephens EH, Borger MA. Getting to the root of the matter: management of the aortic root in type A aortic dissection. *Eur J Cardiothorac Surg.* 2016;50(2):230-231.
8. Tanaka H, Ikeno Y, Abe N, Takahashi H, Inoue T, Okita Y. Outcomes of valve-sparing root replacement in acute type A aortic dissection. *Eur J Cardiothorac Surg.* 2018;53(5):1021-1026.
9. Evangelista A, Isselbacher EM, Bossone E, et al. Insights from the international registry of acute aortic dissection: a 20-year experience of collaborative clinical research. *Circulation.* 2018;137(17):1846-1860.
10. Hess PJ, Jr., Klodell CT, Beaver TM, Martin TD. The Florida sleeve: a new technique for aortic root remodeling with preservation of the aortic valve and sinuses. *Ann Thorac Surg.* 2005;80(2):748-750.
11. Aalaei-Andabili SH, Martin TD, Hess PJ, et al. The Florida sleeve procedure is durable and improves aortic valve function. *Aorta.* 2019;7(2):49-55.
12. Tasca G, Selmi M, Votta E, et al. Aortic root biomechanics after sleeve and david sparing techniques: a finite element analysis. *Ann Thorac Surg.* 2017;103(5):1451-1459.
13. Hess PJ Jr., Harman PK, Klodell CT, et al. Early outcomes using the Florida sleeve repair for correction of aortic insufficiency due to root aneurysms. *Ann Thorac Surg.* 2009;87(4):1161-1168.
14. Mehta RL, Kellum JA, Shah SV, et al. Acute kidney injury network: report of an initiative to improve outcomes in acute kidney injury. *Crit Care.* 2007;11(2):R31.
15. Beckmann E, Martens A, Pertz J, et al. Valve-sparing David I procedure in acute aortic type A dissection: a 20-year experience with more than 100 patients. *Eur J Cardiothorac Surg.* 2017;52(2): 319-324.
16. Subramanian S, Leontyev S, Borger MA, Trommer C, Misfeld M, Mohr FW. Valve-sparing root reconstruction does not compromise survival in acute type A aortic dissection. *Ann Thorac Surg.* 2012; 94(4):1230-1234.
17. Aubin H, Akhyari P, Rellecke P, et al. Valve-sparing aortic root replacement as first-choice strategy in acute type A aortic dissection. *Front Surg.* 2019;6:46.
18. Sultan I, Komlo CM, Bavaria JE, How I. Teach a valve-sparing root replacement. *Ann Thorac Surg.* 2016;101(2):422-425. doi:10.1016/j.athoracsur.2015.12.035
19. Sultan I, McGarvey J, Vallabhajosyula P, Desai ND, Bavaria JE, Szeto WY. Routine use of hemiarach during acute type A aortic dissection repair. *Ann Cardiothorac Surg.* 2016;5(3):245-247. doi:10.21037/acs.2016.04.01
20. Nishimura RA, Otto CM, Bonow RO, et al. AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Thorac Cardiovasc Surg.* 2014; 148(1):e1-e132.

How to cite this article: Alhussaini M, Jeng EI, Martin TD, Fillion A, Beaver TM, Arnaoutakis GJ.. Florida sleeve is a safe and effective technique for valve salvage in acute stanford type A aortic dissection. *J Card Surg.* 2022;37:39-46. doi:10.1111/jocs.16078